

## Adverse Reactions

Very common ( $\geq 1/10$ )	Factor VIII inhibition in previously untreated patients (PUPs)
Common ( $\geq 1/100$ to <1/10)	Headache, Pyrexia
Uncommon ( $\geq 1/1000$ to <1/100)	Influenza, Laryngitis, Factor VIII inhibition in previously treated patients (PTPs), Lymphangitis, Dizziness, Dysgeusia, Memory impairment, Migraine, Syncope, Tremor, Eye inflammation, Palpitations, Haematoma, Hot flush, Pallor, Dyspnoea, Abdominal pain upper, Diarrhoea, Nausea, Vomiting, Hyperhidrosis, Pruritus, Rash, Urticaria, Chest discomfort, Chest pain, Chills, Feeling abnormal, Peripheral oedema, Vessel puncture site haematoma, Coagulation factor VIII level decreased, Haematocrit decreased, Laboratory test abnormal, Monocyte Count increased, Post procedural complication, Post procedural haemorrhage, Procedural site reaction
Not known	Anaphylactic reaction, Hypersensitivity, Fatigue, Injection site reaction, Malaise

## Adverse Reactions

Very common ( $\geq 1/10$ )	Headache
Common ( $\geq 1/100$ to <1/10)	Dizziness, Diarrhoea, Nausea, Rash
Uncommon ( $\geq 1/1000$ to <1/100)	Factor VIII inhibition in previously-treated patients (PTPs), Hypersensitivity, Ocular hyperaemia, Flushing, Drug eruption, Eosinophil count increased, Infusion related reaction